

Change of entropy for one-dimensional ballistic heat equation – sinusoidal initial perturbation

Aleksei A. Sokolov,^{1,2,*} Anton M. Krivtsov,^{2,3,†} Wolfgang H. Müller,^{1,‡} and Elena N. Vilchevskaya^{2,3,§}

¹*Technische Universität Berlin, Einsteinufer 5, 10587 Berlin, Germany*

²*Peter the Great Saint Petersburg Polytechnic University,
Politekhnicheskaja 29, 195251 Saint Petersburg, Russia*

³*Institute for Problems in Mechanical Engineering of the Russian Academy of Sciences,
Bol'shoy pr. 61, V.O., 199178 Saint Petersburg, Russia*

(Dated: July 12, 2018)

This work presents a thermodynamic analysis of the ballistic heat equation from two viewpoints: Classical Irreversible Thermodynamics (CIT) and Extended Irreversible Thermodynamics (EIT). A formula for calculating the entropy within the framework of EIT for the ballistic heat equation is derived. The entropy is calculated for a sinusoidal initial temperature perturbation by using both approaches. The results obtained from CIT show that the entropy is a non-monotonic function and that the entropy production can be negative. The results obtained for EIT show that the entropy is a monotonic function and that the entropy production is nonnegative. A comparison between the entropy behaviors predicted for the ballistic, for the ordinary Fourier-based, and for the hyperbolic heat equation is made. A crucial difference of the asymptotic behavior of the entropy for the ballistic heat equation is shown. It is argued that mathematical time reversibility of the partial differential ballistic heat equation is not consistent with its physical irreversibility. The processes described by the ballistic heat equation are irreversible because of the entropy increase.

I. INTRODUCTION

Classical thermodynamic approaches lead to the Fourier based parabolic law of heat conduction. The propagation of heat described by the classical heat conduction equation is observed on the macroscale and frequently used in engineering applications. However, considering the problem of heat conduction from the atomistic viewpoint can lead to different results. Various models of lattices (considering anharmonicity of the interatomic bonds, 2D lattices) are frequently used for the description of heat transfer [1–15]. Modern technology allowed to demonstrate experimentally [16] that Fourier's law is violated in low-dimensional nanostructures, where a ballistic type of heat conduction is observed. This motivates recent interest in properties of structures, such as graphene and carbon nanotubes [17, 18] and, in particular, in their thermal properties [19]. From a theoretical point of view one of the most attractive playgrounds for the investigation of heat conduction is the harmonic one-dimensional crystal, since all of its thermodynamic properties can be obtained analytically from the equations of lattice dynamics. A pioneering contribution regarding the one-dimensional crystal was made by Schrödinger [20]. He obtained an exact solution for the displacements with arbitrary initial conditions in terms of Bessel functions. This work became a foundation for a future investigation of energy transfer in a one-dimensional chain by Hemmer [21]. A new approach for the description

of non-equilibrium heat conduction processes in crystals was developed in [22–24]. A hyperbolic equation called “ballistic heat equation” was obtained as a mathematical consequence of the equations of lattice dynamics. From an experimental point of view such processes can be observed in low dimensional structures exposed to a laser excitation [25].

The mathematical properties of the ballistic heat equation were investigated in several papers including [26]. The ballistic heat equation is reversible with respect to a substitution of t to $-t$. However, it seems intuitively probable that processes described by the ballistic heat equation are irreversible. The calculation of the entropy production associated with processes described by the ballistic heat equation will help to reveal its thermodynamic properties and answer the question of reversibility. Pioneering ideas considering irreversible thermodynamical processes were mainly developed by Prigogine and Onsager [27, 28]. Their investigations led to the formulation of Classical Irreversible Thermodynamics (CIT). However, further research showed that CIT could not describe a wide class of phenomena including short time and small space scales, as well as hyperbolic models of heat conduction (Cattaneo type) allowing entropy production to be negative in some cases [29]. The phenomenological investigation [30] of the ballistic heat equation showed indeed that within the framework of CIT the entropy production of processes described by the ballistic heat equation can be negative.

One of the modifications of existing theories was Extended Irreversible Thermodynamics (EIT) introduced in [31] and explained in details in [32]. It was shown in [29, 31, 33, 34] that this theory is applicable for more complex models of heat conduction, such as the hyperbolic heat equation (Cattaneo type).

* sokolovalexey1@gmail.com

† akrivtsov@bk.ru

‡ wolfgang.h.mueller@tu-berlin.de

§ vilchevska@gmail.com

The aim of this work is to consider the ballistic heat equation within the framework of EIT, to obtain general formulae, which allow to calculate the entropy for this model, to consider the particular problem of a sinusoidal initial temperature perturbation, and to compare the results obtained by CIT and EIT for the ballistic heat equation, hyperbolic and classical heat conduction equations.

II. MODELS OF HEAT CONDUCTION

If the heat capacity at a constant volume c_V does not depend on time, the energy balance equation reads:

$$\rho c_V \dot{T} = -h', \quad (1)$$

where ρ is the density, T is the temperature and h is the heat flux, and the dot ($\dot{}$) and the dash ($'$) denote temporal and spatial derivatives, respectively. In this paper we are going to analyze three models of heat conduction.

a. The classical heat equation:

$$\dot{T} = \alpha T'', \quad (2)$$

where $\alpha = \kappa/\rho c_V$ is the coefficient of thermal diffusivity, κ is the coefficient of thermal conductivity. Eqn. (2) is obtained on the basis of Fourier's law:

$$h = -\kappa T'. \quad (3)$$

b. Hyperbolic equation (Maxwell-Cattaneo-Vernotte type):

$$\ddot{T} + \frac{1}{\tau} \dot{T} = \frac{\alpha}{\tau} T'', \quad (4)$$

where τ is the relaxation time. It was obtained by introducing a heat flux relaxation term to (3) [29]:

$$\dot{h} + \frac{1}{\tau} h = -\frac{\kappa}{\tau} T'. \quad (5)$$

c. The ballistic heat equation [22]:

$$\ddot{T} + \frac{1}{t} \dot{T} = c^2 T'', \quad (6)$$

where c is the speed of sound, t is the time passed from the moment of the instantaneous heat perturbation. The corresponding equation for the heat flux reads:

$$\dot{h} + \frac{1}{t} h = -\rho k_B c^2 T'. \quad (7)$$

Eqn. (6) was first derived in [22] from the equations of lattice dynamics of a one-dimensional crystal with nearest neighbor linear force interaction:

$$\ddot{q}_i = \omega_e^2 (q_{i-1} - 2q_i + q_{i+1}), \quad \omega_e \stackrel{\text{def}}{=} \sqrt{\frac{C}{m}}, \quad (8)$$

where q_i is the displacement of the particle with index i , C is the interatomic bond stiffness, and m is the particle mass.

The following initial conditions are considered:

$$q_i|_{t=0} = 0, \quad \dot{q}_i|_{t=0} = \sigma(x)\rho_i, \quad (9)$$

where ρ_i are independent random variables with zero expectation and unit variance; σ is the variance of the initial particle velocity. The variance is a slowly changing function of the spatial coordinate $x = ia$, where a is the initial distance between neighboring particles.

The following definitions of the kinetic temperature and the heat flux were used for the derivation:

$$k_B T = m \langle \dot{q}_i \rangle^2, \quad h = \frac{1}{2} C \langle (q_i - q_{i+1})(\dot{q}_i - \dot{q}_{i+1}) \rangle, \quad (10)$$

where $\langle \rangle$ denotes the expectation value. According to the Dulong-Petit law for a one-dimensional system, Boltzmann's coefficient k_B at the right hand side of Eqn. (7) is the specific heat capacity of the one-dimensional crystal at a constant volume c_V .

The classical heat equation Eqn. (2) is solved in combination with the following initial condition:

$$T|_{t=0} = \theta_0(x). \quad (11)$$

Eqs. (4) and (6) are to be solved by using the initial conditions:

$$T|_{t=0} = \theta_0(x), \quad h|_{t=0} = 0. \quad (12)$$

According to (1) the above condition for the heat flux is equivalent to:

$$\dot{T}|_{t=0} = 0. \quad (13)$$

The hyperbolic equation (4) and the ballistic heat equation (12) have similar forms and a somewhat similar behavior (e.g., a finite velocity of the heat front propagation). For the ballistic heat equation (6) the speed of heat propagation is the speed of sound of the medium, $c = \omega_e a$ [22]. For the hyperbolic heat equation (4) the speed of heat propagation is $c_{MCV}^2 = \alpha/\tau$ [35].

However, there is a significant difference between these two equations: the material constant τ is replaced in the ballistic heat equation by the physical time, t [22].

From the form of Eqn. (6) it seems that it has a singularity. However, when Eqn. (6) is solved together with the initial conditions (12), (13) the singularity is absent, which is confirmed by the general analytical solution [36]:

$$T(x, t) = \frac{1}{2\pi} \int_0^{2\pi} T_0(x + ct \cos p) dp \quad (14)$$

and solutions of the particular initial problems [26].

We would like to note that the exact solution for the atomic velocities and displacements (see [20, 21]) of the Eqn. (8) predicts an infinite speed of signal propagation, while the ballistic equation (6) describes a propagation

of temperature at a finite speed, c . This difference is due to the fact that Eqn. (6) is obtained from Eqn. (8) by using continualization and coarse graining in space [22].

Also note that Eqn. (6) describes ballistic heat conduction and does not describe a transition from ballistic to diffusive regimes. It is known that the ballistic heat transport occurs when phonons can propagate without scattering. It happens when the size of the system is comparable to the mean-free path of the carriers. In this case the thermal conductivity is size dependent. However, in a harmonic one-dimensional chain the phonon mean free path is infinite [29] and no phonon-phonon, phonon-impurity, or phonon-boundary scattering occurs. Thus the ballistic heat equation (6) has no size effect and contains only one parameter — the speed of sound in the medium c .

In this paper we will consider and compare the entropy for the models described above, namely: Fourier's heat equation, a phenomenological model which describes heat conduction at the macroscale; the hyperbolic heat equation, a modification of the previous one taking into account wave properties of the heat propagation; and the ballistic heat equation, which is obtained as a direct consequence of the lattice dynamics and is fully based on the wave processes in the crystal lattice.

III. ENTROPY INEQUALITY FOR THE BALLISTIC HEAT EQUATION

The formalism of Classical Irreversible Thermodynamics (CIT) is based on the hypothesis of local equilibrium. It postulates [29] that a thermodynamic system can be divided into a number of microscopic cells, each of which can be treated like a macroscopic system in equilibrium. In each cell the state variables remain uniform but they can change from cell to cell [37]. They can also change with time so that they finally depend continuously on space and time coordinates, (x, t) [29]. Following [30] we consider in this work thermal perturbations only without the presence of mechanical motion. It means that the set of state variables is narrowed down to the specific internal energy, u , only. That leads us to the following form of a *Gibbs relation* [29],

$$ds = \frac{1}{T} du, \quad (15)$$

where s is the specific entropy. However, models taking the independent character of fluxes into account, turn out to be inconsistent with approaches of CIT [29, 30]. Extended irreversible thermodynamics (EIT) introduced a way to avoid contradictions by considering new state variables among the set of basic independent variables. Let us demonstrate this approach when applied to the ballistic heat equation (6). We assume that the entropy depends not only on the internal energy but also on the heat flux, h (in particular the entropy depends on h^2 because

it is independent of the heat flux direction) [29, 31]:

$$s = s(u, h^2) \quad \Rightarrow \quad \dot{s} = \frac{\partial s}{\partial u} \dot{u} + 2 \frac{\partial s}{\partial h^2} \dot{h}. \quad (16)$$

Let us write the Clausius-Duhem inequality in general form [29, 30, 38]:

$$\rho(T\dot{s} - \dot{u}) - \frac{hT'}{T} \geq 0, \quad (17)$$

As shown in [30] the substitution of the second relation in (16) into the inequality (17) yields:

$$\rho \left(T \frac{\partial s}{\partial u} - 1 \right) \dot{u} + h \left(2\rho T \frac{\partial s}{\partial h^2} \dot{h} - \frac{T'}{T} \right) \geq 0, \quad (18)$$

It is assumed in [29, 39] that arbitrary energy supplies keep the balance of energy satisfied. Thus the balance law does not impose constraints on \dot{u} . Therefore in (18) \dot{u} can take arbitrary independent values. Thus in order to guarantee that (18) is satisfied it follows that:

$$\frac{\partial s}{\partial u} = \frac{1}{T}, \quad h \left(2\rho T^2 \frac{\partial s}{\partial h^2} \dot{h} - T' \right) \geq 0. \quad (19)$$

The relations (19) were discussed in previous work [30].

Let us now consider the second inequality in (19) in context with the equation of ballistic thermal conductivity (6). By expressing T' with the first Eqn. from (7) and then substituting it into the inequality from (19) we obtain:

$$\dot{h} h \left(2\rho^2 k_B T^2 c^2 \frac{\partial s}{\partial h^2} + 1 \right) + \frac{h^2}{t} \geq 0. \quad (20)$$

In order to keep this inequality satisfied for any values of \dot{h} , the first term must be zero. Therefore:

$$\frac{\partial s}{\partial h^2} = -\frac{1}{2\rho^2 c^2 c_V T^2}. \quad (21)$$

By substituting the first relation from (19) and the relation (21) into (16) yields:

$$\begin{aligned} \dot{s} &= \frac{1}{T} \dot{u} - \frac{1}{\rho^2 c^2 c_V T^2} h \dot{h} = \dot{s}_{\text{eq}}(u) + \dot{s}_{\text{ne}}(h), \\ \dot{s}_{\text{eq}}(u) &= \frac{1}{T} \dot{u}, \quad \dot{s}_{\text{ne}} = -\frac{1}{\rho^2 c^2 c_V T^2} h \dot{h}, \end{aligned} \quad (22)$$

and as a differential, as an analogy to (15):

$$ds = \frac{1}{T} du - \frac{1}{\rho^2 c^2 c_V T^2} h dh, \quad (23)$$

where s_{eq} is the equilibrium part of the entropy change, which depends only on the specific internal energy, and s_{ne} is the non-equilibrium part, which is dependent on the heat flux.

Thus by considering additional parameters of state (heat flow), one can avoid contradictions leading to a

violation of the second law when using the formulation of CIT. At the same time, the fact that heat can flow from cold to hot, which is observed for the ballistic heat equation (6), is not paradoxical, because it is caused by the inertia of the process under consideration.

Let us suppose that the temperature deviations are small. Then the heat capacity at constant volume, c_V , can be considered to be constant. By postulating further that the internal energy is a function of temperature we obtain:

$$du = c_V dT. \quad (24)$$

By taking into account (24) the relations (15) and (22) have the following differential form:

$$\dot{s}_{\text{CIT}} = c_V \frac{\dot{T}}{T}, \quad (25)$$

$$\dot{s}_{\text{EIT}} = c_V \frac{\dot{T}}{T} - \frac{1}{\rho^2 c^2 c_V T^2} h \dot{h}. \quad (26)$$

The functions describing the temperature $T(x, t)$ and the heat flux $h(x, t)$ are obtained as solutions of the Eqns. (2)–(7). Then by integrating (25) and (26) the corresponding entropies are found.

IV. SINUSOIDAL INITIAL HEAT PERTURBATION

We now present an application of the relations for the entropy in the CIT approach (15) and in the EIT approach (22). To this end consider a sinusoidal initial temperature distribution in an infinite adiabatic one-dimensional system of the form:

$$\theta_0(x) = \delta T \cos kx + T_0, \quad (27)$$

where δT and T_0 are positive constants, which have the dimension of a temperature, and k is a wave number. Let us suppose that functions describing the temporal and spatial evolution of temperature and of the heat flux have the form:

$$\begin{aligned} T(t, x) &= \delta T f_T(\omega t) \cos kx + T_0, \\ h(t, x) &= -\delta T c_V \rho c f_h(\omega t) \sin kx, \end{aligned} \quad (28)$$

where f_T, f_h are dimensionless functions of a dimensionless quantity, ωt , where ω is a parameter with the dimension 1/s. In order to fulfill the initial conditions we put in (28) $f_T(0) = 1, f_h(0) = 0$. Since the perturbation and the solution are laterally periodic we can consider a lateral interval of one period. The wave length of the initial perturbation is $L = 2\pi/k$. We consider the interval $x \in [-L/2, L/2]$. Note that $h(t, -L/2) = h(t, L/2) = 0$, thus there is no heat flow into or out of the system.

In the case of an adiabatically closed system as described above, there is no entropy flux coming in or out

of the system according to [29]. In this case the *rate* of total entropy is equal to the total entropy *production* in the system, which should be non-negative. In this paper we consider an adiabatic system, because in this case a decrease in the total entropy indicates negative entropy production.

A. Classical Irreversible Thermodynamics

We substitute the first equation from (28) into (25) and then perform a series expansion by a small parameter $\frac{\delta T}{T_0}$ (since the temperature deviations are small and $\delta T \ll T_0$) up to terms of second order:

$$\begin{aligned} \dot{s} &= c_V \frac{\delta T}{T_0} \dot{f}_T(\omega t) \cos(kx) \\ &\quad - c_V \left(\frac{\delta T}{T_0} \right)^2 \dot{f}_T(\omega t) f_T(\omega t) \cos^2(kx) + O^3 \left(\frac{\delta T}{T_0} \right). \end{aligned} \quad (29)$$

The total entropy *rate* for the considered interval is given by integration over the whole system:

$$\dot{S}(t) = \int_{-L/2}^{L/2} \rho \dot{s} dx = -\frac{1}{2} c_V \rho L \dot{f}_T(\omega t) f_T(\omega t) \left(\frac{\delta T}{T_0} \right)^2. \quad (30)$$

After integrating over time we obtain:

$$\Delta S_{\text{CIT}}(t) = \frac{1}{4} c_V \rho L (1 - f_T^2(\omega t)) \left(\frac{\delta T}{T_0} \right)^2, \quad (31)$$

where $\Delta S(t) = S(t) - S_0$ is the *change* of entropy, S_0 denotes the initial entropy of the system.

B. Extended Irreversible Thermodynamics

The expression for the entropy change (26) obtained in EIT consists of two terms. One term is the equilibrium part, which is dependent only on the internal energy. It was already calculated above by using the CIT approach. Due to the linearity of integration we can now calculate the non-equilibrium part and then add both results to obtain the full entropy. By substituting the relations (28) into the relation for s_{ne} from (22) and by performing a series expansion with a small parameter $\frac{\delta T}{T_0}$ up to terms of second order we obtain:

$$\dot{s}_{\text{ne}} = -c_V f_h(\omega t) \dot{f}_h(\omega t) \left(\frac{\delta T}{T_0} \right)^2 \cos^2(kx) + O^3 \left(\frac{\delta T}{T_0} \right). \quad (32)$$

The total entropy rate for the considered interval is given by:

$$\dot{S}_{\text{ne}}(t) = \int_{-L/2}^{L/2} \rho \dot{s} dx = -\frac{1}{2} c_V \rho L \dot{f}_h(\omega t) f_h(\omega t) \left(\frac{\delta T}{T_0} \right)^2. \quad (33)$$

After integrating over time we obtain:

$$S_{\text{ne}}(t) = -\frac{1}{4} c_V \rho L f_h^2(\omega t) \left(\frac{\delta T}{T_0} \right)^2. \quad (34)$$

Here the initial non-equilibrium part of the entropy of the system is zero, since there are no fluxes at the beginning: $S_{\text{ne}0} = 0$, so $\Delta S_{\text{ne}} = S_{\text{ne}}(t) - S_{\text{ne}0} = S_{\text{ne}}(t)$. By using this result and (31) and (34) we obtain a relation for the total change of entropy:

$$\Delta S_{\text{EIT}} = \frac{1}{4} c_V \rho L (1 - f_T^2(\omega t) - f_h^2(\omega t)) \left(\frac{\delta T}{T_0} \right)^2. \quad (35)$$

We proceed to specify the functions f_T and f_h , which depend on the chosen model of heat conduction.

V. CALCULATION OF THE ENTROPY FOR DIFFERENT MODELS OF HEAT CONDUCTION

A. Ballistic heat equation

In the case of the ballistic heat equation (6) the solutions for the temperature and for the heat flux have the following forms:

$$\begin{aligned} T(t, x) &= \delta T J_0(kct) \cos kx + T_0, \\ h(t, x) &= \delta T k_B \rho c J_1(kct) \sin kx, \end{aligned} \quad (36)$$

where $J_n(x)$ is the Bessel function of the first kind of order n . By applying the formulae (31) and (35) we have $\omega = kc$, $f_T = J_0$, $f_h = J_1$:

$$\begin{aligned} \Delta S_{\text{CIT}}^{\text{B}} &= \frac{1}{4} c_V \rho L (1 - J_0^2(kct)) \left(\frac{\delta T}{T_0} \right)^2, \\ \Delta S_{\text{EIT}}^{\text{B}} &= \frac{1}{4} c_V \rho L (1 - J_0^2(kct) - J_1^2(kct)) \left(\frac{\delta T}{T_0} \right)^2, \end{aligned} \quad (37)$$

where the superscript B means that entropies corresponds to the ballistic heat equation (6).

B. Hyperbolic equation (Maxwell-Cattaneo-Vernotte)

The entropy for the one-dimensional hyperbolic equation (4), was considered in detail in [33]. Following [33] we represent the results obtained for the entropy pertinent to the hyperbolic heat equation (4) in order to compare them with the ballistic heat equation and the classical heat equation (2). The solution of Eqns. (4) with the initial conditions (12) for a sinusoidal initial perturbation (27) will be the following (see Appendix A):

$$\begin{aligned} 4\alpha\tau k^2 < 1 : \\ T(t, x) &= \delta T e^{-\frac{t}{2\tau}} (\cosh \omega t + A \sinh \omega t) \cos kx + T_0, \\ h(t, x) &= 2c_V \rho \alpha \delta T A e^{-\frac{t}{2\tau}} \sinh \omega t \sin kx, \\ 4\alpha\tau k^2 > 1 : \\ T(t, x) &= \delta T e^{-\frac{t}{2\tau}} (\cos \omega^* t + A^* \sin \omega^* t) \cos kx + T_0, \\ h(t, x) &= 2c_V \rho \alpha \delta T A^* e^{-\frac{t}{2\tau}} \sin \omega^* t \sin kx, \end{aligned} \quad (38)$$

where $\omega = \frac{\sqrt{1-4\alpha\tau k^2}}{2\tau}$, $A = \frac{1}{\sqrt{1-4\alpha\tau k^2}}$, $\omega^* = \frac{\sqrt{4\alpha\tau k^2-1}}{2\tau}$, $A^* = \frac{1}{\sqrt{4\alpha\tau k^2-1}}$, $\kappa = \rho c_V \alpha$ is taken into account. The entropy change reads as follows (from (31) and (35)):

$$\begin{aligned} 4\alpha\tau k^2 < 1 : \\ \Delta S_{\text{CIT}}^{\text{H}} &= \frac{c_V \rho L}{4} \left(1 - e^{-\frac{t}{\tau}} (\cosh \omega t + A \sinh \omega t)^2 \right) \left(\frac{\delta T}{T_0} \right)^2, \\ 4\alpha\tau k^2 > 1 : \\ \Delta S_{\text{CIT}}^{\text{H}} &= \frac{c_V \rho L}{4} \left(1 - e^{-\frac{t}{\tau}} (\cos \omega^* t + A^* \sin \omega^* t)^2 \right) \left(\frac{\delta T}{T_0} \right)^2, \end{aligned} \quad (39)$$

where the superscript H means that entropies corresponds to the hyperbolic heat equation (4). Recall that the formula (22) we use for calculating the entropy based on EIT was obtained for the ballistic heat equation (6). In order to calculate the entropy for the hyperbolic heat equation (4) we use a different formula originally obtained in [29, 31]:

$$\dot{s} = \frac{1}{T} \dot{u} - \frac{\tau}{\rho^2 c_V \alpha T^2} h \dot{h}. \quad (40)$$

This leads to the following relations for entropy:

$$\begin{aligned} 4\alpha\tau k^2 < 1 : \\ \Delta S_{\text{EIT}}^{\text{H}} &= \frac{1}{4} c_V \rho L \left(1 - e^{-\frac{t}{\tau}} \left((\cosh \omega t + A \sinh \omega t)^2 - 4\alpha\tau k^2 A^2 e^{-\frac{t}{\tau}} \sinh^2 \omega t \right) \right) \left(\frac{\delta T}{T_0} \right)^2, \\ 4\alpha\tau k^2 > 1 : \\ \Delta S_{\text{EIT}}^{\text{H}} &= \frac{1}{4} c_V \rho L \left(1 - e^{-\frac{t}{\tau}} \left((\cos \omega^* t + A^* \sin \omega^* t)^2 - 4\alpha\tau k^2 A^{*2} e^{-\frac{t}{\tau}} \sin^2 \omega^* t \right) \right) \left(\frac{\delta T}{T_0} \right)^2. \end{aligned} \quad (41)$$

C. Fourier heat conduction equation.

Let us now consider an application of Eqn. (31) obtained from CIT to the classical Fourier based heat equation (2). The solution of Eqn. (2) for a sinusoidal initial distribution has the form [40]:

$$T(t, x) = \delta T e^{-\alpha k^2 t} \cos kx + T_0. \quad (42)$$

Here we do not consider the heat flux, since CIT does not take it into account as state variable. By applying Eqn. (31) we find with $\omega = -\alpha k^2$, $f_T = e^{-\alpha k^2 t}$:

$$\Delta S_{\text{CIT}}^{\text{F}} = \frac{1}{4} c_V \rho L \left(1 - e^{-2\alpha k^2 t} \right) \left(\frac{\delta T}{T_0} \right)^2, \quad (43)$$

where the superscript F means that the entropy corresponds to the Fourier-based classical heat conduction equation (2).

VI. DIMENSIONAL ANALYSIS

The formulae for entropy obtained above, namely (37), (39), (41), (43) are functions of time with the same dimension. Now we introduce a dimensionless entropy:

$$\Delta\tilde{S} = \frac{4}{\rho L c_V} \left(\frac{T_0}{\delta T} \right)^2 \Delta S. \quad (44)$$

However, dimensionless time can be chosen differently depending on the model of heat conduction. It is well known that the description of physical processes does not depend on the choice of dimensions. Thus it is very useful to describe the process in dimensionless form. The description of heat conduction on different scales leads to different models of heat conduction. However, we would like to compare different properties of these models. Following [41], where the application of *Buckingham's π -theorem* was demonstrated in context with a problem coupling hydrodynamics and electrodynamics, we will apply this theorem to obtain dimensionless parameters, which describe the system above. We will construct and analyze a dimensional matrix A as follows: The basic dimensions of the problem are time - $[T]$, length - $[L]$, mass - $[M]$, temperature - $[\Theta]$. The brackets $[\cdot]$ shows the dimension of a quantity. Then the dimension matrix reads:

$$\begin{array}{c|cccccc} & T & x & t & \alpha & \tau & c \\ \hline [L] & 0 & 1 & 0 & 2 & 0 & 1 \\ [T] & 0 & 0 & 1 & -1 & 1 & -1 \\ [\Theta] & 1 & 0 & 0 & 0 & 0 & 0 \end{array} \implies A_{ij} = \begin{bmatrix} 0 & 1 & 0 & 2 & 0 & 1 \\ 0 & 0 & 1 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}_{ij}. \quad (45)$$

The components of this matrix are exponents of the dimensions. Each column gives the exponents for one considered physical quantity, *i.e.*, $[t] = [L]^0 [T]^1 [\Theta]^0 [M]^0$. The number p of dimensionless quantities Π_i , which is required to describe the system is the number n of physical quantities minus the rank r of the matrix A . In the considered problem we have $n = 6$, $r = 3$. This means that the number p of dimensions is $p = n - r = 3$. They are expressed as a product of physical quantities. $\Pi_i = T^{l_1^i} x^{l_2^i} t^{l_3^i} \alpha^{l_4^i} \tau^{l_5^i} c^{l_6^i}$. As Π_i are dimensionless the following system of linear equations is obtained:

$$\sum_{j=1}^6 A_{ij} l_j^i = 0, \quad i = 1, 2, 3. \quad (46)$$

The exponents l_j^i are the components of the vectors from null space of the matrix A (the null space can be found by solving a system of linear equations (46)). Three linearly independent vectors are given by:

$$\begin{aligned} \mathbf{I}^1 &= [0 \quad -2 \quad 1 \quad 1 \quad 0 \quad 0]; & \mathbf{I}^2 &= [0 \quad 0 \quad -1 \quad 0 \quad 1 \quad 0]; \\ \mathbf{I}^3 &= [0 \quad -1 \quad 1 \quad 0 \quad 0 \quad 1]. \end{aligned} \quad (47)$$

These vectors are needed to construct the following dimensionless quantities:

$$\Pi_1 = \frac{t\alpha}{x^2}, \quad \Pi_2 = \frac{\tau}{t}, \quad \Pi_3 = \frac{ct}{x}. \quad (48)$$

By substituting dimensional coordinates $x = x_{\text{ref}} \tilde{x}$, $t = t_{\text{ref}} \tilde{t}$, $T = T_{\text{ref}} \tilde{T}$ where tildes indicate dimensionless quantities. Then Π_i are redefined by means of reference scales:

$$\Pi_1 = \frac{t_{\text{ref}} \alpha}{x_{\text{ref}}^2}, \quad \Pi_2 = \frac{\tau}{t_{\text{ref}}}, \quad \Pi_3 = \frac{c t_{\text{ref}}}{x_{\text{ref}}}. \quad (49)$$

By using these dimensionless quantities our equations can be rewritten in the dimensionless form (see Table I).

TABLE I: Dimensionless form of heat equations

Ballistic	Hyperbolic	Classical
$\frac{\partial^2 \tilde{T}}{\partial \tilde{t}^2} + \frac{1}{\tilde{t}} \frac{\partial \tilde{T}}{\partial \tilde{t}} = \Pi_3^2 \frac{\partial^2 \tilde{T}}{\partial \tilde{x}^2}$	$\Pi_2 \frac{\partial^2 \tilde{T}}{\partial \tilde{t}^2} + \frac{\partial \tilde{T}}{\partial \tilde{t}} = \Pi_1 \frac{\partial^2 \tilde{T}}{\partial \tilde{x}^2}$	$\frac{\tilde{T}}{\partial \tilde{t}} = \Pi_1 \frac{\partial^2 \tilde{T}}{\partial \tilde{x}^2}$

The reference length scale x_{ref} of our problem is the wavelength of initial periodic perturbation, or we can choose the quantity inverse to wavenumber k : $x_{\text{ref}} = 1/k$. Hence our parameters will be the following ones:

$$\Pi_1 = t_{\text{ref}} \alpha k^2, \quad \Pi_2 = \frac{\tau}{t_{\text{ref}}}, \quad \Pi_3 = c k t_{\text{ref}}. \quad (50)$$

Now an appropriate time scale t_{ref} should be chosen. The reference time scale can be obtained in three different ways: $t_{\text{ref}} = 1/\alpha k^2$ will be interpreted as the time scale of thermal diffusivity leading to $\Pi_1 = 1$ and leaving Π_2 and Π_3 free; $t_{\text{ref}} = \tau$ will be interpreted as the time scale of relaxation time leading to $\Pi_2 = 1$ and leaving Π_1 and Π_3 free; $t_{\text{ref}} = 1/c k$ will be interpreted as the time which a heat wave described by the ballistic heat equation (6) needs to travel along one period of initial perturbation leading to $\Pi_3 = 1$ and leaving Π_1 and Π_2 free.

VII. RESULTS

In this work three heat equations are considered: classical (2), hyperbolic (4), and the ballistic heat equation (6). By using *Buckingham's π theorem*, the system of these three equations can be rewritten in dimensionless form, see Table I. By using the formulae (37), (39), (41), (43) and the results obtained in the previous section we can obtain relations for dimensionless entropy. A comparison of the three models is presented in Table II, where $\tilde{\omega} = \omega \tau$, $\tilde{\omega}^* = \omega^* \tau$. Plots based on all three formulae with the time scale $t_{\text{ref}} = 1/c k$ and the parameters $\Pi_1 = 4$, $\Pi_2 = 1$ are shown in Fig. 1. Figures, similar to Fig. 1 can be found in [33]. They contained the plots for the entropy, which was calculated with CIT and EIT for the hyperbolic heat equation (4). In Fig. 1 we add new results for the ballistic heat equation (6). It is seen that all three equations, the classic (2), the hyperbolic (4), and the ballistic (6) one, lead to same value of dimensionless entropy change at initial time and infinite time, no matter which approach we use, CIT or EIT.

Let us now consider the asymptotic behavior of the obtained entropies. An approximation formula for the Bessel function $J_n(t)$ at large times t ,

TABLE II: Entropy change for sinusoidal initial temperature perturbation.

Equation	Ballistic	Hyperbolic	Classical
Formula	$\ddot{T} + \frac{1}{\tau}\dot{T} = cT''$	$\tau\ddot{T} + \dot{T} = \alpha T''$	$\dot{T} = \alpha T''$
$\Delta\tilde{S}$ CIT	$1 - J_0^2(\tilde{t})$	$4\alpha\tau k^2 < 1 :$ $1 - e^{-\tilde{t}} (A \sinh \tilde{\omega}\tilde{t} + \cosh \tilde{\omega}\tilde{t})^2,$ $4\alpha\tau k^2 > 1 :$ $1 - e^{-\tilde{t}} (A^* \sin \tilde{\omega}^*\tilde{t} + \cos \tilde{\omega}^*\tilde{t})^2$	$1 - e^{-2\tilde{t}}$
$\Delta\tilde{S}$ EIT	$1 - J_0^2(\tilde{t}) - J_1^2(\tilde{t})$	$4\alpha\tau k^2 < 1 :$ $1 - e^{-\tilde{t}} (A \sinh \tilde{\omega}\tilde{t} + \cosh \tilde{\omega}\tilde{t})^2 - 4\alpha\tau k^2 C^2 \sinh^2 \tilde{\omega}\tilde{t},$ $4\alpha\tau k^2 > 1 :$ $1 - e^{-\tilde{t}} (A^* \sin \tilde{\omega}^*\tilde{t} + \cos \tilde{\omega}^*\tilde{t})^2 - 4\alpha\tau k^2 C^{*2} \sin^2 \tilde{\omega}^*\tilde{t},$	—
Time scale	$\tilde{t} = kct$	$\tilde{t} = \frac{t}{\tau}$	$\tilde{t} = \alpha k^2 t$
Asymptotics	$1/\tilde{t}$	$e^{-\tilde{t}}$	$e^{-\tilde{t}}$




FIG. 1: Plots of the entropy for ballistic heat equation calculated using CIT for hyperbolic (dashed red line), CIT for ballistic heat equation (dashed black line), EIT for hyperbolic (solid red line), EIT for ballistic equation (solid black line) and CIT for classical heat equation (solid blue line).

is given by the following expression [42]:

$$J_n(t) \approx \sqrt{\frac{2}{\pi t}} \cos\left(t - \frac{n\pi}{2} - \frac{\pi}{4}\right) + \mathcal{O}\left(\frac{1}{t}\right). \quad (51)$$

According to (51) the relations for the entropy for the ballistic heat equation from (37) reach the asymptotic value as a power law. The formulae (41) and (43) indicate that the entropy for the hyperbolic (4) and classical (2) heat conduction will asymptotically be exponential. Also note that entropy values at initial time 0 and at large times $t \rightarrow \infty$ are given by 0 and 1 as seen from Fig. 1.

VIII. CONCLUSIONS

In this work three equations of heat conduction were investigated, the classical (2), the hyperbolic (4), and the ballistic one (6). The ballistic equation (6) of heat conduction was not considered from a phenomenological thermodynamics point of view before. This paper shows that the CIT approach can lead to a negative entropy production for the ballistic heat equation (6). Therefore the ballistic heat equation (6) is considered within the framework of EIT. A general formula (22) is obtained for calculating the entropy for Eqn.(6) with an arbitrary initial temperature perturbation. The example of an adiabatically closed system with an initial sinusoidal temperature perturbation is considered. For such a system a non-monotonic increase of the total entropy indicates negative entropy production. This example shows that the CIT approach is not applicable for the ballistic equation (6) and causes a negative entropy production and a non-monotonic “wavy” increase of the entropy. The entropy calculated with formula (35) obtained for EIT increases monotonically, and the entropy production stays non-negative.

We would like to note that in the case of CIT only the temperature (as a function of x and t) is needed to calculate the entropy (see Eqn. (25)). The expression for the temperature for a wide class of scalar lattices was obtained in [24]. Thus the results obtained in the current work within the framework of CIT can be extended to the case of scalar lattices.

Regarding the asymptotic behavior of entropy the following can be said: There are papers showing that a harmonic system consisting of an infinite number of particles will approach spatial equilibrium for large times, see for example [43]. It is seen that at time $t \rightarrow \infty$, *i.e.*, when the system tends to equilibrium, the value of entropy is equal for all three models, no matter which approach is used, EIT or CIT. The change of dimensionless entropy $\Delta\tilde{S}$ tends to one at large times. However, the asymptotic behavior of the entropy is significantly

different. The entropy calculated for hyperbolic and for classical Fourier equations tends exponentially, e^{-t} , to an asymptotic value, whereas the entropy for ballistic heat equation (6) behaves according to a power law, $1/t$.

If we now consider the problem from a discrete point of view, which in full generality is beyond the scope of this paper, then it can be said that the harmonic crystal can be decomposed into a system of independent modes and there will be no energy exchange between them. But in this system the irreversibility and the entropy rise is associated with the phases of oscillations, which distribute independently for long times — the details can be found in the pioneering work by P.C. Hemmer [21].

We want to conclude by mentioning a frequently accepted concept of reversibility: Obviously the equation of ballistic heat conduction (6)

$$\ddot{T} + \frac{1}{t}\dot{T} = c^2 T'' \quad (52)$$

is invariant with respect to time reversion: $t \rightarrow -t$. However, the results presented above show the irreversible nature of the ballistic heat conduction (6) process, because of the increase of the total entropy of the system. Thus the mathematical time reversibility of the ballistic PDE (6) is not correlated with its physical irreversibility.

ACKNOWLEDGMENTS

This work is supported by joint grant of the German Research Foundation (DFG) (No. 405631704) and the Russian Science Foundation (No. 19-41-04106). Contribution of W.H. Müller and A.A. Sokolov is supported by the German Research Foundation, contribution of A.M. Krivtsov and E.N. Vilchevskaya is supported by the Russian Science Foundation.

The authors of this work would like to thank Dmitry Korikov and Vitaly Kuzkin for the useful discussions.

Appendix A: Solution for hyperbolic heat equation with sinusoidal initial temperature perturbation

In this section we present the solution of the hyperbolic heat equation (4) with the initial conditions (27). We find the solution in the form

$$T(t, x) = \delta T f(t) \cos kx + T_0. \quad (A1)$$

Substitution of (A1) leads us to the following ODE for $f(t)$,

$$\tau \ddot{f} + \dot{f} + \alpha k^2 f = 0, \quad (A2)$$

with the following initial conditions:

$$f(0) = 1, \quad \dot{f}(0) = 0. \quad (A3)$$

The solution is found in the form e^{pt} . The corresponding characteristic equation for the Eqn. (A2) is:

$$\tau p^2 + p + \alpha k^2 = 0. \quad (A4)$$

Depending on the ratio of the parameters α , k , τ this equation has real or complex roots:

$$\begin{aligned} 4\alpha\tau k^2 < 1 : \quad p_{1,2} &= \frac{-1 \pm \sqrt{1-4\alpha\tau k^2}}{2\tau}, \\ 4\alpha\tau k^2 > 1 : \quad p_{1,2} &= \frac{-1 \pm i\sqrt{4\alpha\tau k^2-1}}{2\tau}. \end{aligned} \quad (A5)$$

It leads us to the following fundamental solutions for Eqn. (A2):

$$\begin{aligned} 4\alpha\tau k^2 < 1 : \\ f(t) &= A_1 e^{-\frac{1+\sqrt{1-4\alpha\tau k^2}}{2\tau}t} + A_2 e^{-\frac{1-\sqrt{1-4\alpha\tau k^2}}{2\tau}t}, \\ 4\alpha\tau k^2 > 1 : \\ f(t) &= e^{-\frac{t}{2\tau}} \left(B_1 \cos \frac{\sqrt{-1+4\alpha\tau k^2}}{2\tau}t + B_2 \sin \frac{\sqrt{-1+4\alpha\tau k^2}}{2\tau}t \right). \end{aligned} \quad (A6)$$

The coefficients A_1 , A_2 , B_1 , B_2 are found by substitution of Eqns. (A6) into the initial conditions (A3):

$$\begin{aligned} A_1 &= \frac{1+\sqrt{1-4\alpha\tau k^2}}{\sqrt{1-4\alpha\tau k^2}}, \quad A_2 = \frac{-1+\sqrt{1-4\alpha\tau k^2}}{\sqrt{1-4\alpha\tau k^2}}, \\ B_1 &= 1, \quad B_2 = \frac{1}{\sqrt{-1+4\alpha\tau k^2}}. \end{aligned} \quad (A7)$$

α is the coefficient of thermal diffusivity. By using representations of hyperbolic trigonometric functions the solution for the temperature is:

$$\begin{aligned} 4\alpha\tau k^2 < 1 : \\ T(t, x) &= \delta T e^{-\frac{t}{2\tau}} (\cosh \omega t + A \sinh \omega t) \cos kx + T_0, \\ 4\alpha\tau k^2 > 1 : \\ T(t, x) &= \delta T e^{-\frac{t}{2\tau}} (\cos \omega^* t + A^* \sin \omega^* t) \cos kx + T_0. \end{aligned} \quad (A8)$$

The solutions for the heat flux are obtained by substitution of Eqns. (A8) into the second Eqn. from (4):

$$\begin{aligned} 4\alpha\tau k^2 < 1 : \\ \tau \dot{h} + h &= \alpha \rho c_V k \delta T e^{-\frac{t}{2\tau}} (\cosh \omega t + A \sinh \omega t) \sin kx, \\ 4\alpha\tau k^2 > 1 : \\ \tau \dot{h} + h &= \alpha \rho c_V k \delta T e^{-\frac{t}{2\tau}} (\cos \omega^* t + A^* \sin \omega^* t) \sin kx. \end{aligned} \quad (A9)$$

The solution for the Eqns. (A9) is found in the form:

$$h(t) = \alpha \rho c_V k \delta T f_h(t) \sin kx. \quad (A10)$$

This leads us to the following inhomogeneous ODE's:

$$\begin{aligned} 4\alpha\tau k^2 < 1 : \tau \dot{f}_h + f_h &= e^{-\frac{t}{2\tau}} (\cosh \omega t + A \sinh \omega t), \\ 4\alpha\tau k^2 > 1 : \tau \dot{f}_h + f_h &= e^{-\frac{t}{2\tau}} (\cos \omega^* t + A^* \sin \omega^* t), \end{aligned} \quad (A11)$$

with the initial condition

$$f_h(0) = 0. \quad (A12)$$

The fundamental solutions for Eqns. (A11) are:

$$\begin{aligned} 4\alpha\tau k^2 < 1 : f_h &= D_1 e^{-\frac{t}{\tau}} + 2A e^{-\frac{t}{2\tau}} \sinh \omega t, \\ 4\alpha\tau k^2 > 1 : f_h &= D_2 e^{-\frac{t}{\tau}} + 2A^* e^{-\frac{t}{2\tau}} \sin \omega^* t. \end{aligned} \quad (A13)$$

By substituting Eqns. (A11) into the initial conditions (A12) we obtain:

$$D_1 = 0, \quad D_2 = 0. \quad (A14)$$

The solution for the heat flux is then:

$$\begin{aligned} 4\alpha\tau k^2 < 1 : h(t, x) &= 2c_V\rho\alpha\delta T A e^{-\frac{x}{2\tau}} \sinh \omega t \sin kx, \\ 4\alpha\tau k^2 > 1 : h(t, x) &= 2c_V\rho\alpha\delta T A^* e^{-\frac{x}{2\tau}} \sin \omega^* t \sin kx. \end{aligned} \quad (\text{A15})$$

-
- [1] A. Dhar and R. Dandekar, *Physica A: Statistical Mechanics and its Applications* **418**, 49 (2015), proceedings of the 13th International Summer School on Fundamental Problems in Statistical Physics.
- [2] A. V. Savin and Y. A. Kosevich, *Phys. Rev. E* **89**, 032102 (2014).
- [3] M. B. Babenkov, A. M. Krivtsov, and D. V. Tsvetkov, *Physical Mesomechanics* **19**, 282 (2016).
- [4] D. Saadatmand, D. Xiong, V. A. Kuzkin, A. M. Krivtsov, A. V. Savin, and S. V. Dmitriev, *Phys. Rev. E* **97**, 022217 (2018).
- [5] S. Lepri, R. Livi, and A. Politi, *Physics Reports* **377**, 1 (2003).
- [6] S. N. Gavrilov, A. M. Krivtsov, and D. V. Tsvetkov, *Continuum Mechanics and Thermodynamics* (2018).
- [7] R. Goldstein and N. Morozov, *Physical Mesomechanics* **10**, 235 (2007).
- [8] M. A. Guzev, *Dal'nevost. Mat. Zh.* **18**, 34 (2018).
- [9] M. C. Zheng, F. M. Ellis, T. Kottos, R. Fleischmann, T. Geisel, and T. Prosen, *Phys. Rev. E* **84**, 021119 (2011).
- [10] G. Falasco, M. Baiesi, L. Molinaro, L. Conti, and F. Baldovin, *Phys. Rev. E* **92**, 022129 (2015).
- [11] D. Roy and A. Dhar, *Phys. Rev. E* **78**, 051112 (2008).
- [12] A. V. Savin, G. P. Tsironis, and X. Zotos, *Phys. Rev. B* **75**, 214305 (2007).
- [13] G. T. Landi and M. J. de Oliveira, *Phys. Rev. E* **87**, 052126 (2013).
- [14] H. Zhou, G. Zhang, J.-S. Wang, and Y.-W. Zhang, *Phys. Rev. E* **94**, 052123 (2016).
- [15] V. Kannan, A. Dhar, and J. L. Lebowitz, *Phys. Rev. E* **85**, 041118 (2012).
- [16] C. W. Chang, D. Okawa, H. Garcia, A. Majumdar, and A. Zettl, *Phys. Rev. Lett.* **101**, 075903 (2008).
- [17] A. Porubov and I. Berinskii, *International Journal of Non-Linear Mechanics* **67**, 27 (2014).
- [18] I. Berinskii, A. Krivtsov, and A. Kudarova, *Physical Mesomechanics* **17**, 356 (2014).
- [19] C. Faugeras, B. Faugeras, M. Orlita, M. Potemski, R. Nair, and A. Geim, *ACS Nano* **4**, 1889 (2010).
- [20] E. Schrödinger, *Annalen der Physik* **44**, 916 (1914).
- [21] P. C. Hemmer, *Dynamic and stochastic types of motion in linear chain*, Ph.D. thesis, Norwegian Institute of Technology (1959).
- [22] A. M. Krivtsov, *Doklady Physics* **60**, 407 (2015).
- [23] A. M. Krivtsov, *Doklady Physics* **59**, 427 (2014).
- [24] V. A. Kuzkin and A. M. Krivtsov, *J. Phys.: Condens. Matter* **29**, 505401 (2017).
- [25] D. A. Indeitsev and E. V. Osipova, *Doklady Physics* **62**, 136 (2017).
- [26] A. A. Sokolov, A. M. Krivtsov, and W. H. Müller, *Physical Mesomechanics* **20**, 305 (2016).
- [27] I. Prigogine, *Introduction to thermodynamics of irreversible processes* (New York, Interscience Publishers, 1961).
- [28] L. Onsager, *Phys. Rev.* **37** (1931).
- [29] D. Jou, G. Lebon, and J. Casas-Vázquez, *Extended Irreversible Thermodynamics* (Springer, Netherlands, 2010).
- [30] A. M. Krivtsov, A. A. Sokolov, W. H. Müller, and A. Freidin, "Advances in mechanics of microstructured media and structures," (Springer, Cham, 2017) Chap. One-dimensional heat conduction and entropy production.
- [31] I. Müller, *Zur Ausbreitungsgeschwindigkeit von Störungen in kontinuierlichen Medien*, Ph.D. thesis, RWTH Aachen (1966).
- [32] I. Müller and T. Ruggeri, *Extended Thermodynamics* (Springer-Verlag, New York, 1993).
- [33] M. Criado-Sancho and J. E. Llebot, *Physical Review E* **47**, 4104 (1993).
- [34] F. X. Alvarez, J. Casas-Vázquez, and D. Jou, *Phys. Rev. E* **77**, 031110 (2008).
- [35] M. Özişik and B. Vick, *International Journal of Heat and Mass Transfer* **27**, 1845 (1984).
- [36] A. M. Krivtsov and V. A. Kuzkin, "Encyclopedia of continuum mechanics," (Springer-Verlag GmbH Germany, 2018) Chap. Discrete and Continuum Thermomechanics.
- [37] S. R. De Groot and P. Mazur, *Non-Equilibrium Thermodynamics* (North-Holland Publishing Company, 1962).
- [38] V. Palmov, *Vibrations of Elasto-Plastic Bodies* (Springer-Verlag Berlin Heidelberg, 1998).
- [39] B. Coleman, *Arch Rat Mech Anal* **17**, 1 (1964).
- [40] H. S. Carslaw and J. C. Jaeger, *Conduction of Heat in Solids* (Clarendon Press, Oxford., 1959).
- [41] S. Glane, F. A. Reich, and W. H. Müller, *Continuum Mech. Thermodyn.* , 1 (2018).
- [42] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions: With Formulas, Graphs, and Mathematical Tables* (Dover Publications, 1964).
- [43] H. Spohn and J. L. Lebowitz, *Communications in Mathematical Physics* **54**, 97 (1977).